

# EVC/Bioregion Benchmark for Vegetation Quality Assessment

## Gippsland Plains bioregion

### EVC 14: Banksia Woodland

#### Description:

Open scrub to open woodland to 10 m tall on well-drained Quaternary and Tertiary sandy soils. The overstorey is dominated by Saw Banksia *Banksia serrata*, sometimes with emergent eucalypts while the understorey is shrub-rich.

#### Large trees:

Species	DBH(cm)	#/ha
<i>Banksia serrata</i>	50 cm	15 / ha

#### Tree Canopy Cover:

%cover	Character Species	Common Name
15%	<i>Banksia serrata</i>	Saw Banksia

#### Understorey:

Life form	#Spp	%Cover	LF code
Immature Canopy Tree		5%	IT
Understorey Tree or Large Shrub	1	5%	T
Medium Shrub	10	25%	MS
Small Shrub	7	15%	SS
Large Herb	1	1%	LH
Medium Herb	5	10%	MH
Small or Prostrate Herb	1	1%	SH
Large Tufted Graminoid	3	10%	LTG
Large Non-tufted Graminoid	1	5%	LNG
Medium to Small Tufted Graminoid	5	10%	MTG
Medium to Tiny Non-tufted Graminoid	2	5%	MNG
Ground Fern	2	10%	GF
Scrambler or Climber	3	1%	SC
Bryophytes/Lichens	na	10%	BL

# EVC 14: Banksia Woodland - Gippsland Plains bioregion

LF Code	Species typical of at least part of EVC range	Common Name
MS	<i>Epacris impressa</i>	Common Heath
MS	<i>Leptospermum continentale</i>	Prickly Tea-tree
MS	<i>Acacia terminalis</i>	Sunshine Wattle
MS	<i>Acacia suaveolens</i>	Sweet Wattle
SS	<i>Dampiera stricta</i>	Blue Dampiera
SS	<i>Amperea xiphoclada</i> var. <i>xiphoclada</i>	Broom Spurge
SS	<i>Tetralochea pilosa</i>	Hairy Pink-bells
SS	<i>Dillwynia glaberrima</i>	Smooth Parrot-pea
MH	<i>Gonocarpus teucrioides</i> s.l.	Germander Raspwort
MH	<i>Selaginella uliginosa</i>	Swamp Selaginella
MH	<i>Drosera peltata</i> ssp. <i>auriculata</i>	Tall Sundew
LTG	<i>Lomandra longifolia</i>	Spiny-headed Mat-rush
LTG	<i>Xanthorrhoea resinosa</i>	Spear Grass-tree
LTG	<i>Gahnia clarkei</i>	Tall Saw-sedge
LNG	<i>Tetrarrhena juncea</i>	Forest Wire-grass
MTG	<i>Dianella caerulea</i> var. <i>caerulea</i>	Paroo Lily
MTG	<i>Patersonia glabrata</i>	Leafy Purple-flag
MTG	<i>Lepidosperma laterale</i>	Variable Sword-sedge
MTG	<i>Burchardia umbellata</i>	Milkmaids
MNG	<i>Empodisma minus</i>	Spreading Rope-rush
GF	<i>Pteridium esculentum</i>	Austral Bracken
GF	<i>Lindsaea linearis</i>	Screw Fern
SC	<i>Cassytha glabella</i>	Slender Dodder-laurel
SC	<i>Cassytha phaeolasia</i>	Rusty Dodder-laurel
SC	<i>Billardiera scandens</i>	Common Apple-berry

## Recruitment:

Episodic/Fire. Desirable period between disturbances is 30 years.

## Organic Litter:

40 % cover

## Logs:

15 m/0.1 ha.

## Weediness:

There are no consistent weeds in this EVC.

Published by the Victorian Government Department of Sustainability and Environment November 2007

© The State of Victoria Department of Sustainability and Environment 2007

This publication is copyright. Reproduction and the making available of this material for personal, in-house or non-commercial purposes is authorised, on condition that:

- the copyright owner is acknowledged;
- no official connection is claimed;
- the material is made available without charge or at cost; and
- the material is not subject to inaccurate, misleading or derogatory treatment.

Requests for permission to reproduce or communicate this material in any way not permitted by this licence (or by the fair dealing provisions of the *Copyright Act 1968*) should be directed to the Nominated Officer, Copyright, 8 Nicholson Street, East Melbourne, Victoria, 3002.

For more information contact: Customer Service Centre, 136 186

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

[www.dse.vic.gov.au](http://www.dse.vic.gov.au)